

General Topology and its Applications 9 (1976) 117–125
 © North-Holland Publishing Company

PARACOMPACTNESS AND PRODUCTS

Scott W. WILLIAMS

*Department of Mathematics, State University of New York at Buffalo, Amherst,
 N.Y. 14226, U.S.A.*

Received 7 May 1974

We investigate paracompactness in the product of a paracompact space Y with a paracompact linearly ordered topological space X , where certain controls are placed on the subspace ηX of points without a compact neighborhood. If ηX is dispersed, then $X^\alpha \times Y$ is paracompact for all finite ordinals α . If the Lindelöf degree of Y is less than the least cardinal number, for each $n \in \eta X$, of any collection of neighborhoods whose intersection does not contain n in its interior, then $X \times Y$ is paracompact.

AMS Subj. Class.: Primary 54D20; Secondary 54F05

paracompact	dispersed
product spaces	scattered
linearly ordered topological spaces	

Many of the standard examples showing that the product of two paracompact spaces is not always paracompact require one or both factors to be a linearly ordered topological space (LOTS) or a subspace of a LOTS. We illustrate some sufficient conditions for the product of two paracompact spaces to be paracompact when one factor is a LOTS (or a subspace of a LOTS). All spaces are assumed Hausdorff.

1. Notation

A *linearly ordered topological space* (LOTS) is a linearly ordered set with its interval topology. An *interior gap* of a LOTS X is a Dedekind cut $(A|B)$ of X such that A has no supremum (sup) and B has no infimum (inf). An *end-gap*, left or right, means the absence of an infimum

(inf) or supremum (sup) of the linearly ordered set. The *Dedekind compactification* X^+ of a LOTS X is formed by suitably ordering $X \cup \{g: g \text{ is a gap of } X\}$ in a manner similar to the completion of the rationals. X is dense in the compact LOTS X^+ . A gap is a *Q-gap* provided it is the limit of an increasing sequence, if not a right end-gap, and a decreasing sequence, if not a left end-gap, each having no limit points in X . For further details on LOTS, their gaps, and their compactifications, we suggest [4].

If X is any linearly ordered set, then X^* denotes the same set with the inverse order of X .

Intervals in a LOTS X are denoted by $[a, b]$ when closed and by $]a, b[$ when open, and in the latter case a and/or b may be allowed to be a gap. A convex set C satisfies $a, b \in C$ and $a < x < b$ implies $x \in C$. Singleton sets are considered to be convex.

Let X be a LOTS, for $x \in X^+$ let $\tau_+(x)$ (resp. $\tau_-(x)$) be the smallest ordinal λ for which there is a decreasing (resp. increasing) sequence of type λ in X^+ coinitial (resp. cofinal) with x . Then the *type* of x is $\tau(x) = \inf\{\tau_+(x), \tau_-(x)\}$ while the *character* of x is $\chi(x) = \sup\{\tau_+(x), \tau_-(x)\}$.

The *Lindelöf degree* $L(X)$ of a topological space is the smallest cardinal α such that each open covering of X has a subcovering of cardinal less than or equal to α .

For a topological space X , we define ηX to be the subspace of all points of X without a compact neighborhood.

A topological space is called *dispersed*¹ if every non-empty subspace has an isolated point.

In this paper we make a slight deviation from the usual definition for the term "refinement" in that a refinement of a covering need not be a covering of any set unless it is so stated.

2. Remarks and examples

A list of papers concerning the normality or paracompactness of products would be long, and hence we shall not present one here; however, certain of these results should be mentioned for their strong relationship with our work.

2.1. Katuta [6] has shown that if X is paracompact, and $X \times Y$ is normal for every paracompact space Y , then $X \times Y$ is paracompact.

¹ Dispersed spaces are also called scattered spaces in the literature.

2.2. The disjoint union of the Michael line [8] and the Sorgenfrey line [10] can be embedded as a closed subspace of a first-countable hereditarily paracompact LOTS L using a technique of Fedorčuk [3] or Lutzer (see discussion after Corollary 3.3). Hence $L \times L$ and $L \times P$ are not normal, where P is the space of irrationals.

2.3. Przymusiński [9], armed with Martin's Axiom and $2^{\aleph_0} = 2^{\aleph_1}$, has given the Sorgenfrey topology to a subset X of \mathbb{R} to produce a hereditarily paracompact space whose product with itself is perfectly normal and not paracompact.

2.4. That a separable LOTS is paracompact follows from results in [4]. That a separable LOTS is also an M-space² is known as a "folk result" to the author [13]. It is well-known that the product of two paracompact M-spaces is paracompact.

2.5. Each of the spaces L in 2.2 and X in 2.3 is a paracompact non-M-space. Consider the space Y to be the long line [11] with the last point added, the countable ordinals subtracted, and given the interval topology. Y is a paracompact non-M-space; however, Theorem 3.3 will show that $Y \times Z$ is paracompact for any Lindelöf space Z . Further, Theorem 3.4 will show that $Y \times Z$ is paracompact for any paracompact space Z . On the other hand, the results in [12] do not seem to be applicable in this example.

2.6. For results concerning normality in products of spaces, we suggest [1], [2] and [5].

3. Theorems

As a trivial corollary of a result of Suzuki [12] we obtain a very necessary lemma. Each of our theorems is a generalization of this result.

3.1. Lemma. *If X and Y are paracompact and X is a locally compact LOTS, then $X \times Y$ is paracompact.*

² X is an M-space if there is a closed continuous function f from X onto a metric space Y such that $f^{-1}(y)$ is countably compact for each $y \in Y$.

3.2. Lemma [4]. *A LOTS X is paracompact if and only if every gap of X is a Q-gap.*

3.3. Theorem. *Let X and Y be paracompact spaces, and let X be a LOTS for which $\tau(n) > L(Y)$ or $\tau(n) = 0$ and $\chi(n) > L(Y)$ for each $n \in \eta X$. Then $X \times Y$ is paracompact.*

Proof. Let R be an open covering of $X \times Y$ and $A \subseteq X$. For convenience we say that S is a *pro-cover* of A provided S is an open σ -locally-finite refinement of R covering $A \times Y$.

For $x_1, x_2 \in X$ define $x_1 \sim x_2$ if and only if there is a pro-cover of

$$[\inf \{x_1, x_2\}, \sup \{x_1, x_2\}] .$$

Then \sim is an equivalence relation on X whose classes we designate $kl(x)$. It will be shown that x is an interior point of $kl(x)$ and there is a pro-cover of $kl(x)$ for each $x \in X$. From this it follows that $X \times Y$ is paracompact.

x is an interior point of $kl(x)$. Indeed, it is obvious from Lemma 3.1 if $x \notin \eta X$. On the other hand, if $x \in \eta X$, there is a refinement R_1 of R covering $\{x\} \times Y$ such that R_1 consists entirely of members each of the form $C \times U$, where C is open and convex in X and U is open in Y . Choose an open locally-finite refinement H of $\{U: C \times U \in R_1\}$ covering Y , and for each $W \in H$ choose $C(W)$ to be open and convex in X and $U(W)$ open in Y such that $W \subseteq U(W)$ and $C(W) \times U(W) \in R_1$. Then

$$R_2 = \{C(W) \times W: W \in H\}$$

is a pro-cover of $\{x\}$. Let $R_3 \subseteq R_2$ be a covering of $\{x\} \times Y$ such that $|R_3| \leq L(Y)$.

Suppose $\tau_+(x) > 0$; then $\tau_+(x) > |R_3|$, so

$$x < \inf \{\sup C(W): C(W) \times W \in R_3\} .$$

Similarly, $\tau_-(x) > 0$ implies

$$x > \sup \{\inf C(W): C(W) \times W \in R_3\} .$$

In any case, x is in the interior of

$$\cap \{C(W): C(W) \times W \in R_3\},$$

and R_3 is a pro-cover of the latter. Therefore, x is an interior point of $\text{kl}(x)$; therefore, $\text{kl}(x)$ is an open and closed convex set in X .

Suppose $\inf(\text{kl}(x)) \in X$ and $\sup(\text{kl}(x)) = g \notin X$, then $\inf(\text{kl}(x))$ is in the interior of $\text{kl}(\inf(\text{kl}(x)))$, from which it follows that $\inf(\text{kl}(x)) \in \text{kl}(x)$ and $\inf(\text{kl}(x)) = \inf X$. So there is a pro-cover T of $[\inf X, x]$. On the other hand, from Lemma 3.2, there is an increasing sequence

$$\{x_\beta: \beta < \tau_-(g)\} \subseteq X^+$$

cofinal with g such that

- (1) $x_0 = x$,
- (2) if $\tau_-(g) = \omega$, we may choose $x_\beta \in X$ for all β ,
- (3) $\{x_\beta: \beta < \tau_-(g)\}$ has no limit points in X .

If $\tau_-(g) = \omega$, we choose S_β to be a pro-cover of $[x_0, x_{\beta+1}]$ for all $\beta < \omega$. In this case,

$$T \cup \cup \{S_\beta: \beta < \omega\}$$

is a pro-cover of $\text{kl}(x)$.

If $\tau_-(g) > \omega$, we choose S_0 to be a pro-cover of $[x_0, x_1[$ and S_β to be a pro-cover of

$$] \sup \{x_\gamma: \gamma < \beta\}, x_{\beta+1} [$$

for all $\beta > 0$ such that

$$\cup \{V: V \in S_\beta\} =] \sup \{x_\gamma: \gamma < \beta\}, x_{\beta+1} [\times Y.$$

It is clear that each of the following is a σ -locally-finite collection:

- (a) $(S_0 \cup T) \cap ([\inf X, x_1[\times Y)$;
- (b) $\cup \{S_\lambda: \lambda \text{ is a limit ordinal}\}$;
- (c) $\{V: V \in S_{\lambda+k}, \lambda \text{ is a limit ordinal}, 0 < \lambda < \tau_-(g)\}$ for $0 < k < \omega$.

Therefore, the union of these collections is a pro-cover of $\text{kl}(x)$.

There are three other cases to examine: $\inf(\text{kl}(x)) \in X$ and $\sup(\text{kl}(x)) \in X$; $\inf(\text{kl}(x)) \notin X$ and $\sup(\text{kl}(x)) \in X$; $\inf(\text{kl}(x)) \notin X$ and $\sup(\text{kl}(x)) \notin X$. However, in each case a pro-cover of $\text{kl}(x)$ can be constructed with a modification of the techniques of the previous paragraphs.

Since $\text{kl}(x)$ is clopen for each $x \in X$ and $\text{kl}(x)$ has a pro-cover P_x for each $x \in X$ such that

$$\bigcup P_x = \text{kl}(x) \times Y,$$

it follows that X has a pro-cover. \square

If, in Theorem 3.3, Y is in fact a LOTs, then $\chi(y) \leq L(Y)$ for each $y \in Y^+ - Y$. We conjecture that substitution of " $\tau(n) > \chi(y)$ or $\tau(n) = 0$ and $\chi(n) > \chi(y)$ for each $n \in \eta X$ and $y \in Y^+ - Y$ " is not sufficient to reach $X \times Y$ paracompact.

3.4. Theorem. *Let X and Y be paracompact, and let X be a LOTs for which ηX is dispersed. Then $X \times Y$ is paracompact.*

Proof. For each pair $x_1, x_2 \in X$ we say $x_1 \sim x_2$ if and only if

$$[\inf\{x_1, x_2\}, \sup\{x_1, x_2\}] \times Y$$

is paracompact. Then \sim is an equivalence relation on X whose classes we denote $\text{kl}(x)$. We show that $\text{kl}(x) \times Y$ is paracompact and x is an interior point of $\text{kl}(x)$ for each $x \in X$. From this it follows that $X \times Y$ is paracompact.

Form the derivatives of ηX by allowing $D_0 = \eta X$, D_α to be the set of non-isolated points of $D_{\alpha-1}$ when α is non-limit, and $D_\alpha = \bigcap \{D_\beta : \beta < \alpha\}$ when α is a limit ordinal. For $x \in X$ call $\alpha(x)$ the first ordinal such that $x \notin D_{\alpha(x)}$. Our proof is by induction.

Suppose $\alpha(x) = 0$; we may use Lemma 3.1 to show that x is an interior point of $\text{kl}(x)$. Moreover, if $\sup(\text{kl}(x)) \notin D_0$ or if $\sup(\text{kl}(x))$ is not the limit of an increasing sequence of gaps, then a modification of the third paragraph of Theorem 3.3 shows $[x, \sup(\text{kl}(x))] \times Y$ to be paracompact. Hence we suppose for $k = \sup(\text{kl}(x))$ that $k \in D_0$ and k is the limit of an increasing sequence of gaps and R is an open cover of $[x, k] \times Y$. There exists a collection $R_1 = \{U_i : i \in I\}$ of open sets of $X \times Y$ such that

- (1) each U_i is of the form $C_i \times G_i$ for each $i \in I$,
- (2) C_i is a convex set with $g_i = \inf C_i \notin X$ for each $i \in I$,
- (3) $\{G_i : i \in I\}$ is a locally finite covering of Y ,
- (4) R_1 is a locally finite refinement of R covering $\{k\} \times Y$.

For each $i \in I$ choose S_i to be an open locally finite refinement of R covering $]x, g_i[\times Y$ and let

$$T_i = \{V \cap (]x, g_i[\times G_i) : V \in S_i\}.$$

Then $\{R_1\} \cup \bigcup \{T_i; i \in I\}$ is an open locally finite refinement of R covering $]x, k] \times Y$. Repetition of these techniques shows that $[\inf(\text{kl}(x)), x] \times Y$, and hence $\text{kl}(x) \times Y$, is paracompact and closed in X .

Suppose $n \in \eta X$ and $\alpha(x) < \alpha(n)$ implies $\text{kl}(x) \times Y$ is paracompact, x is an interior point of $\text{kl}(x)$, and $\text{kl}(x)$ is closed. We consider the case $\tau(n) > 0$. There exist $a, b \in X$ such that $n \in]a, b[,]a, n[\neq \emptyset,]n, b[\neq \emptyset, \alpha(a) < \alpha(x), \alpha(b) < \alpha(x)$, and

$$[a, b] \cap D_{\alpha(n)-1} = \{n\}.$$

Whether or not $[a, b]$ is the disjoint union of clopen convex sets each of whose product with Y is paracompact, we have

$$\inf(\text{kl}(b)) \leq n \leq \sup(\text{kl}(a)),$$

since $\text{kl}(c) \times Y$ is paracompact for all $c \in [a, b] - \{n\}$. Since $\text{kl}(a)$ and $\text{kl}(b)$ are closed, $[a, b] \times Y$ is paracompact and n is in the interior of $\text{kl}(n)$. On the other hand, $n \in \text{kl}(a)$, so $\text{kl}(n) \times Y$ is paracompact. The cases for $\tau(n) = 0$ are similar. \square

3.5. Corollary. *If X is a paracompact LOTS and ηX is dispersed, then X^α is paracompact for all finite ordinals α .*

Lutzer [7] has shown that if X is a subspace of a LOTS, then there exists an order-preserving homeomorphism from X onto a closed subspace of a certain LOTS L obtained by replacing points which are

- (1) suprema of sequences but not limit points of those sequences with ω^* ,
- (2) infima of sequences but not limit points of those sequences with ω ,
- (3) suprema and infima of sequences but are isolated points with $\omega^* + \omega$.

Lutzer has shown that L is paracompact whenever X is paracompact. It is easy to show that ηL is dispersed whenever ηX is dispersed. Hence we would have the following:

3.6. Corollary. *If X is a paracompact subspace of a LOTS and ηX is dispersed, then $X \times Y$ is paracompact for any paracompact space Y .*

Similarly, we may apply Theorem 3.3 to arrive at:

3.7. Corollary. *Let X be a paracompact subspace of a LOTS, and let Y be a topological space. If $L(Y)$ is less than the least cardinal number, for each $n \in \eta X$, of any collection of neighborhoods of n whose intersection does not contain n in its interior, then $X \times Y$ is paracompact.*

We make two conjectures sparked by recent theorems and counterexamples in the area of box topologies:³

3.8. Assume the continuum hypothesis. If X_n is a paracompact LOTS for which ηX_n is dispersed for each $n \in \omega$, then $\prod \{X_n : n \in \omega\}$ is paracompact when given the box topology.

3.9. There is a paracompact space X for which ηX is dispersed and a metric space Y such that $X \times Y$ is not paracompact.

The author wishes to express a gratitude to the referee whose suggestions for modification of the original draft of this paper were most helpful.

References

- [1] M. Atsuji, Necessary and sufficient conditions for the normality of the product of two spaces, *Proc. Japan Acad.* 45 (1969) 894–898.
- [2] R. Conover, Normality and products of linearly ordered spaces, *Gen. Topology Appl.* 2 (1972) 215–225.
- [3] V. Fedorčuk, Ordered sets and the products of topological spaces, *Proc. Moscow Univ.* 4 (1966).
- [4] L. Gilman and M. Henriksen, Concerning rings of continuous functions, *Trans. Am. Math. Soc.* 77 (1954) 340–362.
- [5] I. Hayashi, The normality of the product of two linearly ordered spaces, *Proc. Japan Acad.* 43 (1969) 300–304.
- [6] Y. Katuta, On the normality of the product of a normal space with a paracompact space, *Gen. Topology Appl.* 1 (1971) 295–319.
- [7] D.J. Lutzer, On generalized ordered spaces, Ph.D. Thesis, University of Washington, Seattle, Wash. (1970).
- [8] E. Michael, The product of a normal space and a metric space need not be normal, *Bull. Am. Math. Soc.* 69 (1963) 375–376.
- [9] T. Przymusiński, A Lindelöf space X such that X^2 is normal but not paracompact, *Fund. Math.* 78 (1973) 291–296.
- [10] R. Sorgenfrey, On the topological product of paracompact spaces, *Bull. Am. Math. Soc.* 53 (1947) 631–632.

³ The most comprehensive collection of recent results in box topologies known to the author at this time is in a series of notes from lectures given at the University of Wyoming in August 1974 by M.E. Rudin.

- [11] L.A. Steen and J.A. Seebach, Counter-examples in Topology (Holt, Reinhart, and Winston, New York, 1970).
- [12] J. Suzuki, Paracompactness of product spaces, *Proc. Japan Acad.* 45 (1969) 457–460.
- [13] S. Williams, A simple fact of rationals, *Notices Am. Math. Soc. Abstract* 711-54-36, A-218.